

Shingles



Herpes zoster

What is shingles?

Shingles, also known as herpes zoster, is a disease caused by the varicella-zoster virus. This virus also causes the common childhood disease chickenpox. After a person has chickenpox, the varicella-zoster virus lies dormant in their nerves. For unknown reasons, the virus may reactivate later in life, and shingles will appear.

Who gets shingles?

Anyone who has had chickenpox can get shingles. Shingles is more common in persons with depressed immune systems and in persons over the age of 50, but it can also occur in children. Typically the symptoms in children are milder than in adults. People with weakened immune systems may have more severe disease.

How is shingles spread?

Adults and children with shingles are contagious and can transmit the varicella-zoster virus to people who have not had chickenpox or been vaccinated against it.

Direct contact with the rash or broken blisters can transmit the virus, so the rash should be completely covered with a bandage until it crusts over. If the rash cannot be covered, the ill person should stay at home until it crusts over. Children and adults who have not had chickenpox should avoid direct contact with shingles lesions.

What are the symptoms of the shingles?

Shingles typically starts with a burning sensation in a patch of skin, followed by a cluster of small red bumps. This rash will change to painful fluid-filled blisters, which eventually dry and crust over. The rash usually goes away in one to two weeks, but pain may continue for a prolonged period of time. Shingles typically follows the path of a single nerve and occurs on one side of the body. The blisters are commonly found on the torso and buttocks, but shingles may appear on the arms, legs or face.

Health care providers diagnose shingles through physical examination, gentle skin scrapings, and/or blood tests, since shingles sometimes resembles other skin conditions. Your health care provider will determine whether or not to use medication (an antiviral) to treat shingles. Immediately consult a health care provider if the rash involves the eyes,

gets red and drains pus, becomes very itchy and extremely painful, or if the rash lasts more than two weeks.

Is there a vaccine for shingles?

A licensed vaccine is available that reduces the risk of shingles. The vaccine, Zostavax, is FDA approved (3/24/11) for use in those aged 50 years and older.

For more information:

Spokane Regional Health District
Disease Prevention and Response Division
(509) 324-1442 | TDD (509) 324-1464